

# The Greenhouse Effect, a Comparison of Greenhouse Gas Contributions from Fossil Fueled and Geothermal Power Plants with Potential Solutions for Emission Reductions

Leslie Morrison<sup>i</sup>, Brian Benn, Allen Sonnevile<sup>i</sup>, Ken McIntush<sup>ii</sup>  
Calpine Corporation, The Geysers, Sonoma and Lake Counties, CA<sup>i</sup>  
Trimeric Corporation, Buda TX<sup>ii</sup>

## Keywords

*Climate change, global warming, greenhouse effect, greenhouse gas, global warming potential, CO<sub>2</sub> equivalent, Geysers*

## ABSTRACT


Fossil fueled power plants across the world emit greenhouse gases that contribute to global warming. Geothermal power is considered a “green” alternative to coal, oil and gas fired power production. But how green is it? Analysis of the non-condensable gases that come out of The Geysers’ reservoir with the geothermal steam show varying amounts of carbon dioxide and methane. These constituents pass through the power production cycle; the CO<sub>2</sub> is emitted to the atmosphere, and so is the CH<sub>4</sub> unless it is burned as part of the hydrogen sulfide abatement process. This paper reviews a basic understanding of the greenhouse effect, and evaluates the relative contribution of greenhouse gas emissions from a geothermal power plant compared to other forms of power production, and proposes possible solutions to reduce these emissions.

## 1. Introduction

Climate change includes both global warming (the gradual increase in average surface temperature of the Earth) and its impact on Earth’s weather patterns. During Earth’s long history, there have been previous periods of climate change, however, most scientists agree that the current rapid rate of change is due to human-made emissions rather than due to natural phenomena. The so called “Greenhouse Effect” is caused by the emission of gases that trap radiant energy that normally would flow from the Earth’s surface out into space.

Any warm mass radiates energy proportional to its temperature – the Sun at about 5,500 °C (9,930 °F) emits most of its energy as ultraviolet, visible and near infrared light. Approximately 75% of this energy arriving from our Sun passes through Earth’s atmosphere, which does not absorb energy well at these wavelengths, and warms its surface. (25% of the sun’s energy is reflected by Earth’s atmosphere back into space). Earth’s average surface temperature of about

15 °C (59 °F) is much cooler than the sun so it emits longer wavelength infrared radiant heat which is absorbed by natural “greenhouse gases” in our atmosphere. The atmosphere re-radiates energy both upwards and downwards; the part radiated downwards is absorbed again by the surface of Earth. This leads to a higher equilibrium temperature than if the atmosphere did not both absorb and radiate energy. Without the natural greenhouse effect, Earth's average temperature would be well below freezing. Current human-caused increases in greenhouse gases cause greater amounts of energy to be absorbed by the atmosphere and re-radiated by the atmosphere back to Earth, causing the Earth to grow warmer over time.<sup>1</sup>



**Figure 1: Diagram of radiant energy flow between the Sun, Earth, Earth's Atmosphere, and Space.<sup>2</sup>**

Greenhouse gases are specific airborne compounds that can absorb energy in the infrared radiation wavelength. The gases are ranked based on their global warming potential (GWP) which is the heat absorbed by any greenhouse gas in the atmosphere, as a multiple of the heat that would be absorbed by the same mass as the reference gas, carbon dioxide (CO<sub>2</sub>). Thus it provides a common scale for measuring the climate effects of different gases. GWP is 1 for CO<sub>2</sub>.

The GWP depends on the following factors:

- The absorption capacity of infrared radiation (dependent on chemical structure and bonds). A gas has the most effect if it absorbs in a "window" of wavelengths where the absorption by the natural atmospheric components is fairly transparent.
- The time horizon of interest (most regulators use 100 years).
- The atmospheric lifetime of the gas (how quickly the gas species is removed from the atmosphere)<sup>3</sup>.


**Table 1: Global Warming Potential values for CO<sub>2</sub> and CH<sub>4</sub>. Table amended to show only the two most predominant greenhouse gases emitted from geothermal and fossil fueled power plants<sup>4</sup>**

Industrial designation or common name	Chemical formula	GWP values for 100-year time horizon		
		Second Assessment Report (SAR)	Fourth Assessment Report (AR4)	Fifth Assessment Report (AR5)
Carbon dioxide	CO <sub>2</sub>	1	1	1
Methane	CH <sub>4</sub>	21	25	28


Water is actually the strongest greenhouse gas, because it has a profound infrared absorption spectrum with more and broader absorption bands than CO<sub>2</sub>. However, it is removed via precipitation within weeks, so its GWP is negligible.<sup>5</sup>

The major constituents of Earth's atmosphere, nitrogen (N<sub>2</sub> at 78% by volume), oxygen (O<sub>2</sub> at 21%) and argon (Ar at 0.9%), are not greenhouse gases because molecules containing two atoms of the same element, such as N<sub>2</sub> and O<sub>2</sub>, have no net change in the distribution of their electrical charges when they vibrate. Monatomic gases, such as Ar, do not have vibrational modes. Therefore, none of these major constituents can absorb the infrared radiation emitted by the Earth.

In 2020, methane (CH<sub>4</sub>) accounted for about 11% of all U.S. greenhouse gas emissions from human activities. Human activities emitting methane include leaks from natural gas systems and livestock. Methane is also emitted by natural sources such as wetlands. However, natural processes in soil and chemical reactions in the atmosphere help remove CH<sub>4</sub> from the atmosphere. Methane's lifetime in the atmosphere is much shorter than carbon dioxide (CO<sub>2</sub>), but CH<sub>4</sub> is more efficient at trapping radiation than CO<sub>2</sub>. Pound for pound, the comparative impact of CH<sub>4</sub> is 25 times greater than CO<sub>2</sub> over a 100-year period.<sup>6</sup>



Carbon dioxide (CO<sub>2</sub>) is the primary greenhouse gas emitted through human activities. In 2020, CO<sub>2</sub> accounted for about 79% of all U.S. greenhouse gas emissions from human activities. Carbon dioxide is naturally present in the atmosphere as part of the Earth's carbon cycle (the natural circulation of carbon among the atmosphere, oceans, soil, plants, and animals). Human activities are altering the carbon cycle—both by adding more CO<sub>2</sub> to the atmosphere and by reducing the ability of natural sinks, like forests and soils, to remove and store CO<sub>2</sub> from the atmosphere. While CO<sub>2</sub> emissions come from a variety of natural sources, human-related emissions are implicated for the increase that has occurred in the atmosphere since the industrial revolution. CO<sub>2</sub> emissions cause increases in atmospheric concentrations of CO<sub>2</sub> that will last thousands of years.<sup>7</sup>



**Figure 2: Atmospheric Carbon Dioxide (1960-2021)<sup>7</sup>**

Carbon dioxide equivalent (CO<sub>2</sub>e or CO<sub>2</sub>eq or CO<sub>2</sub>-e) is calculated from GWP. For any gas, it is the mass of CO<sub>2</sub> that would warm the earth as much as 1 kg of that gas. Thus, it provides a common scale for measuring the climate effects of different gases. It is calculated as GWP times mass of the other gas. For example, if a gas has GWP of 100, two tonnes of the gas have CO<sub>2</sub>e of 200 tonnes.

## 2. Greenhouse Gas Emissions from Fossil Fueled Power Plants

In 2020, power plants that burned coal, natural gas, and petroleum, accounted for 62% of all U.S. electricity generation, but they account for 99% of U.S. electricity related CO<sub>2</sub> emissions. And even though fossil fueled power plants are being shut down all over the country in preference for cleaner sources of energy such as wind and solar, there is still a need for electricity generation that is not limited by nighttime darkness and calm, windless days.

**Table 2: U.S. Electric Power Generation and CO<sub>2</sub> Emissions by Fuel 2020<sup>8</sup>**

	<b>Electric Generation (MWh)</b>	<b>CO<sub>2</sub> GWP Emissions (Metric Tons)</b>	<b>kg GWP/MWh</b>
<b>Coal</b>	757,763,000	767,000,000	1,012
<b>Natural Gas</b>	1,402,438,000	576,000,000	411
<b>Petroleum</b>	13,665,000	13,000,000	951

(GWP =global warming potential)  
 Combined cycle plants are excluded because some of their CO<sub>2</sub> emissions are from fuel consumption for heating purposes.  
 \*US Energy Information Administration, 2020<sup>8</sup>

## 3. Greenhouse Gas Emissions at The Geysers

Geothermal power is a reliable energy source that is available 24 hours a day, all year round. However, it is not 100% carbon free when it comes to greenhouse gas emissions and global warming potential. But the greenhouse gas emissions from geothermal power are more than an order of magnitude lower than those emitted from fossil fuel fired power plants.

**Table 3: Summary of 2021 Green House Gas Emissions from Calpine Geysers Power Plants<sup>9</sup>**

<b>Net Generation MWh</b>	<b>5,519,164</b>		
<b>Global Warming Potential (GWP) (CO<sub>2</sub> equivalent)</b>	<b>Metric Tons</b>		<b>Metric Tons</b>
	<b>CO<sub>2</sub></b>	<b>CH<sub>4</sub></b>	<b>GWP</b>
	<b>1</b>	<b>25</b>	<b>CO<sub>2</sub> Equivalent</b>
<b>Power plants</b>	126,788	3,021	202,314
			<b>kg GWP/MWh</b>
			37

The steam emanating from The Geysers’ deep underground geothermal reservoir contains less than 1%wt non-condensable gases, including some CO<sub>2</sub>, CH<sub>4</sub> and Hydrogen Sulfide (H<sub>2</sub>S). The exact amounts of these three constituents varies over the geographical area of the known geothermal resource. It is believed that the CO<sub>2</sub>, CH<sub>4</sub>, and H<sub>2</sub>S are natural remnants from volcanic activity in this region over 1.1 million years ago. (which, by the way, is also the source of the heat for the hot, dry steam that makes geothermal power in this area possible.)

Currently, The Geysers only has air pollution control regulatory limits for H<sub>2</sub>S, SO<sub>x</sub>, NO<sub>x</sub>, CO, HAPS>10 TPY and particulate emissions. Of these, only H<sub>2</sub>S and SO<sub>2</sub> require special removal equipment to meet the emission limits. There are several methods used for removing H<sub>2</sub>S from the non-condensable gases that are discharged from the power plant condenser. The thermal oxidation (burner) method at four of the power plants affects the greenhouse gas emissions by converting the CH<sub>4</sub> to CO<sub>2</sub> which is emitted to the atmosphere along with the rest of the CO<sub>2</sub>. This reduces the global warming potential of the CH<sub>4</sub> portion of the NCG emissions by a ratio of 2.75:25. (remember that we need to take into account the fact that newly formed CO<sub>2</sub> (MW = 44) weighs more than CH<sub>4</sub> (MW = 16), so we don’t get the full benefit of 1:25 reduction in global warming potential.)

However, eight of the power plants at The Geysers do not use burners for H<sub>2</sub>S emission control. At these units the CH<sub>4</sub> along with most of the CO<sub>2</sub> passes through the power plants and are released to the atmosphere.

#### 4. Case Study: Lake View-Unit 17

The steam that supplies Unit 17 comes from a relatively high non-condensable gas region of the geothermal reservoir. This unit uses a Stretford process for H<sub>2</sub>S abatement rather than a burner. Consequently, since both the CO<sub>2</sub> and CH<sub>4</sub> will pass through a Stretford process, this power plant has the highest greenhouse gas emissions at the Geysers.

**Table 4: 2021 Green House Gas Emissions from Calpine’s Lake View-Unit 17<sup>9</sup>**

<b>Net Generation MWh</b>	<b>471,638</b>			
	<b>Metric Tons</b>		<b>Metric Tons</b>	<b>Kg GWP/MWh</b>
	<b>CO<sub>2</sub></b>	<b>CH<sub>4</sub></b>	<b>GWP</b>	
<b>Global Warming Potential (GWP) (CO<sub>2</sub> equivalent)</b>	<b>1</b>	<b>25</b>	<b>CO<sub>2</sub> Equivalent</b>	
Unit 17	21,272	809	41,497	88

Although Lake View-Unit 17 emissions have a global warming potential far below the values created by fossil fuel power plants, the question arises, can we do better?

Listed below are several concepts that have been proposed for reducing CO<sub>2</sub> and CH<sub>4</sub> emissions. If the current level of GHG emissions from the geothermal industry are ever deemed unattractive for power purchase contracts, subsidies or tax credits, it may be prudent to evaluate the feasibility and economics of some of these proposals for future use in geothermal power plants.

- 1) Use a waste heat boiler (WHB)<sup>10</sup> to combust any non-CO<sub>2</sub> greenhouse gases (e.g., methane) to less GWP intensive CO<sub>2</sub>. This may provide the extra benefit of generating some additional steam which could
  - a) Supplement the geothermal steam and lower the GWP/MWh.
  - b) Be used in a separate stand-alone power cycle to produce energy which could be classified and valued as “zero emission” power.
- 2) Use CO<sub>2</sub> in greenhouses (possibly after burning the NCG stream in a WHB and recovering heat?)
- 3) Direct injection of NCG underground into a suitable zone (assumes this is possible and does not cause other problems)
- 4) Purification of CO<sub>2</sub> (e.g., to ~90+%) and then injection into a suitable zone purely for sequestration
- 5) Purification of CO<sub>2</sub> to EOR grade and transport it to a location that can use it for enhanced oil recovery (EOR)
- 6) Purification of CO<sub>2</sub> to beverage grade (this CO<sub>2</sub> ends up emitted when the beverage is drunk, but it might displace other CO<sub>2</sub> and may reduce carbon footprint)
- 7) Conversion of CO<sub>2</sub> to methanol, MeOH
- 8) Conversion of CO<sub>2</sub> to other hydrocarbons or chemicals (via syngas route)
- 9) Purification of CO<sub>2</sub> and use as a concrete additive (note, this is a sequestration measure and not a value-adder for concrete. This may have the benefit of qualifying for tax credits)

## 5. Conclusion

Although geothermal power may not be 100% carbon free, its greenhouse gas emissions are far below the amount emitted from conventional fossil fueled power plants on a per megawatt basis. This fact coupled with the potential to lower GHGs even more with enhanced technology, should make geothermal power a preferred choice within the electric industry. The even lower global warming potential could be a valuable selling point for geothermal power.

## 6. References

<sup>1</sup>Rebecca, Lindsay (14 January, 2009). “Climate and Earth’s Energy, Budget Feature Articles”. Earthobservatory.nasa.gov. Archived from the original on 21 January 2021. Retrieved 14 December, 2020

<sup>2</sup>A loose necktie, July 12, 2019 File: Greenhouse-effect-t2.svg Wikimedia Commons, Licensed under the Creative Commons Attribution-Share Alike 4.0 International license

- <sup>3</sup> Matthew Elrod, "Greenhouse Warming Potential Model." Based on Elrod, M. J. (1999). "Greenhouse Warming Potentials from the Infrared Spectroscopy of Atmospheric Gases". Journal of Chemical Education. 76 (12): 1702. Bibcode:1999JChEd..76.1702E. doi:10.1021/ed076p1702.
- <sup>4</sup> Myhre, G., D. Shindell, F.-M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.-F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura and H. Zhang, 2013: *Anthropogenic and Natural Radiative Forcing. In: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA
- <sup>5</sup> Steven C Sherwood<sup>1</sup>, Vishal Dixit and Chryséis Salomez<sup>2</sup> "The Global Warming Potential of Near-Surface Emitted Water Vapor" Published 27 September 2018 • © 2018 The Author(s). Published by IOP Publishing Ltd Environmental Research Letters, Volume 13, Number 10 Citation Steven C Sherwood et al 2018 *Environ. Res. Lett.* **13** 104006
- <sup>6</sup> IPCC (2007). Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. [S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press. Cambridge, United Kingdom 996 pp.
- <sup>7</sup> IPCC (2013). Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. [Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P. M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1585 pp.<sup>7</sup> <https://www.climate.gov/news-features/understanding-climate/climate-change-atmospheric-carbon-dioxide>
- <sup>8</sup> U.S. Energy Information Administration; <https://www.eia.gov/tools/faqs/faq.php?id=74&t=11>
- <sup>9</sup> Sonnevile, Allen, "Summary of Greenhouse Gas Emissions, 2021" Calpine, The Geysers
- <sup>10</sup> Avcry, J. et al. "Use of Waste Heat Boiler to Capture Energy from Flammable Non-Condensable Gas at geothermal Power Plants." GRC Transactions, Vol. 39, 2015